arXiv:1305.1089v2 [math.CO] 14 May 2013

Graphs with large generalized (edge-)connectivity*

Xueliang Li[1], Yaping Mao[1], Lusheng Wang[2]

[1]Center for Combinatorics and LPMC-TJKLC, Nankai University

Tianjin 300071, China. Email: lxl@nankai.edu.cn; maoyaping@ymail.com

[2]Department of Computer Science, City University of Hong Kong

Tat Chee Avenue, Kowloon, Hong Kong SAR, China. Email: lwang@cs.cityu.edu.hk

Abstract

The generalized k-connectivity $\kappa_k(G)$ of a graph G, introduced by Chartrand et al. in 1984, is a nice generalization of the classical connectivity. Recently, as a natural counterpart, we proposed the concept of generalized k-edge-connectivity $\lambda_k(G)$. In this paper, graphs of order n such that $\kappa_k(G) = n - \lceil \frac{k}{2} \rceil - 1$ and $\lambda_k(G) = n - \lceil \frac{k}{2} \rceil - 1$ for even k are characterized, respectively.

Keywords: (edge-)connectivity; Steiner tree; internally disjoint trees; edge-disjoint trees; generalized (edge-)connectivity.

AMS subject classification 2010: 05C40, 05C05, 05C75.

1 Introduction

All graphs considered in this paper are undirected, finite and simple. We refer to the book [1] for graph theoretical notation and terminology not described here. The generalized connectivity of a graph G, introduced by Chartrand et al. in [2], is a natural and nice generalization of the concept of (vertex-)connectivity. For a graph $G = (V, E)$ and a set $S \subseteq V(G)$ of at least two vertices, *an S-Steiner tree* or *a Steiner tree connecting S* (or simply, *an S-tree*) is a such subgraph $T = (V', E')$ of G that is a tree with $S \subseteq V'$. Two Steiner trees T and T' connecting S are said to be *internally disjoint* if $E(T) \cap E(T') = \varnothing$ and $V(T) \cap V(T') = S$. For $S \subseteq V(G)$ and $|S| \geq 2$, the *generalized local connectivity* $\kappa(S)$ is the maximum number of internally disjoint trees connecting S in G. Note that when $|S| = 2$ a Steiner tree connecting S is just a path connecting the two vertices of S. For an integer k with $2 \leq k \leq n$, the *generalized k-connectivity* $\kappa_k(G)$ of G is defined as

---

*Supported by NSFC No.11071130, and RGCHK Project No. CityU 9041382.

$\kappa_k(G) = min\{\kappa(S) : S \subseteq V(G)\ and\ |S| = k\}$. Clearly, when $|S| = 2$, $\kappa_2(G)$ is nothing new but the connectivity $\kappa(G)$ of G, that is, $\kappa_2(G) = \kappa(G)$, which is the reason why one addresses $\kappa_k(G)$ as the generalized connectivity of G. By convention, for a connected graph G with less than k vertices, we set $\kappa_k(G) = 1$. Set $\kappa_k(G) = 0$ when G is disconnected. Results on the generalized connectivity can be found in [3, 10, 11, 12, 13, 14, 17].

As a natural counterpart of the generalized connectivity, we introduced the concept of generalized edge-connectivity in [15]. For $S \subseteq V(G)$ and $|S| \geq 2$, the *generalized local edge-connectivity* $\lambda(S)$ is the maximum number of edge-disjoint trees connecting S in G. For an integer k with $2 \leq k \leq n$, the *generalized k-edge-connectivity* $\lambda_k(G)$ of G is then defined as $\lambda_k(G) = min\{\lambda(S) : S \subseteq V(G)\ and\ |S| = k\}$. It is also clear that when $|S| = 2$, $\lambda_2(G)$ is nothing new but the standard edge-connectivity $\lambda(G)$ of G, that is, $\lambda_2(G) = \lambda(G)$, which is the reason why we address $\lambda_k(G)$ as the generalized edge-connectivity of G. Also set $\lambda_k(G) = 0$ when G is disconnected.

In addition to being natural combinatorial measures, the Steiner Tree Packing Problem and the generalized edge-connectivity can be motivated by their interesting interpretation in practice as well as theoretical consideration.

From a theoretical perspective, both extremes of this problem are fundamental theorems in combinatorics. One extreme of the problem is when we have two terminals. In this case internally (edge-)disjoint trees are just internally (edge-)disjoint paths between the two terminals, and so the problem becomes the well-known Menger theorem. The other extreme is when all the vertices are terminals. In this case internally disjoint trees and edge-disjoint trees are just spanning trees of the graph, and so the problem becomes the classical Nash-Williams-Tutte theorem.

Theorem 1. *(Nash-Williams [16], Tutte [19]) A multigraph G contains a system of ℓ edge-disjoint spanning trees if and only if*

$$\|G/\mathscr{P}\| \geq \ell(|\mathscr{P}| - 1)$$

holds for every partition $\mathscr{P}$ of $V(G)$, where $\|G/\mathscr{P}\|$ denotes the number of crossing edges in G, i.e., edges between distinct parts of $\mathscr{P}$.

The generalized edge-connectivity is related to an important problem, which is called the *Steiner Tree Packing Problem*. For a given graph G and $S \subseteq V(G)$, this problem asks to find a set of maximum number of edge-disjoint Steiner trees connecting S in G. One can see that the Steiner Tree Packing Problem studies local properties of graphs, but the generalized edge-connectivity focuses on global properties of graphs. The generalized edge-connectivity and the Steiner Tree Packing Problem have applications in $VLSI$ circuit design, see [4, 5, 18]. In this application, a Steiner tree is needed to share an electronic signal by a set of terminal nodes. Another application, which is our primary focus, arises in the Internet Domain. Imagine that a given graph G represents a network. We choose arbitrary k vertices as nodes. Suppose that one of the nodes in G is a *broadcaster*, and all the other nodes are either *users* or *routers* (also called *switches*). The broadcaster wants to broadcast as many streams of movies as possible, so that the users have the maximum number of choices. Each stream of movie is broadcasted via a tree connecting all the users and the broadcaster. So, in essence we need to find the maximum number Steiner trees connecting all the users and the broadcaster, namely, we want to get $\lambda(S)$, where S is the set of the k nodes. Clearly, it is a Steiner tree packing problem. Furthermore, if we want to know whether for any k nodes the network G has the above properties, then we need to compute $\lambda_k(G) = min\{\lambda(S)\}$ in order to prescribe the reliability and the security of the network.

The following two observations are easily seen from the definitions.

Observation 1. *Let k, n be two integers with $3 \leq k \leq n$. For a connected graph G of order n, $\kappa_k(G) \leq \lambda_k(G) \leq \delta(G)$.*

Observation 2. *Let k, n be two integers with $3 \leq k \leq n$. If H is a spanning subgraph of G of order n, then $\lambda_k(H) \leq \lambda_k(G)$.*

Chartrand et al. in [2] got the exact value of the generalized k-connectivity for the complete graph K_n.

Lemma 1. *[2] For every two integers n and k with $2 \leq k \leq n$, $\kappa_k(K_n) = n - \lceil k/2 \rceil$.*

In [15] we obtained some results on the generalized k-edge-connectivity. The following results are restated, which will be used later.

Lemma 2. *[15] For every two integers n and k with $2 \leq k \leq n$, $\lambda_k(K_n) = n - \lceil k/2 \rceil$.*

Lemma 3. *[15] Let k, n be two integers with $3 \leq k \leq n$. For a connected graph G of order n, $1 \leq \kappa_k(G) \leq \lambda_k(G) \leq n - \lceil k/2 \rceil$. Moreover, the upper and lower bounds are sharp.*

We also characterized graphs attaining the upper bound and obtained the following result.

Lemma 4. *[15] Let k, n be two integers with $3 \leq k \leq n$. For a connected graph G of order n, $\kappa_k(G) = n - \lceil \frac{k}{2} \rceil$ or $\lambda_k(G) = n - \lceil \frac{k}{2} \rceil$ if and only if $G = K_n$ for even k; $G = K_n \setminus M$ for odd k, where M is an edge set such that $0 \leq |M| \leq \frac{k-1}{2}$.*

Like in [15], here we continue to characterizing the graphs with large generalized k-connectivity and generalized k-edge-connectivity. Graphs of order n such that $\kappa_k(G) = n - \lceil \frac{k}{2} \rceil - 1$ and $\lambda_k(G) = n - \lceil \frac{k}{2} \rceil - 1$ for any even k are characterized, respectively.

Theorem 2. *Let n and k be two integers such that k is even and $4 \leq k \leq n$, and G be a connected graph of order n. Then $\kappa_k(G) = n - \frac{k}{2} - 1$ if and only if $G = K_n \setminus M$ where M is an edge set such that $1 \leq \Delta(K_n[M]) \leq \frac{k}{2}$ and $1 \leq |M| \leq k - 1$.*

The above result can also be established for the generalized k-edge-connectivity, which is stated as follows.

Theorem 3. *Let n and k be two integers such that k is even and $4 \le k \le n$, and G be a connected graph of order n. Then $\lambda_k(G) = n - \frac{k}{2} - 1$ if and only if $G = K_n \setminus M$ where M is an edge set satisfying one of the following conditions:*

(1) $\Delta(K_n[M]) = 1$ *and* $1 \le |M| \le \lfloor \frac{n}{2} \rfloor$*;*

(2) $2 \le \Delta(K_n[M]) \le \frac{k}{2}$ *and* $1 \le |M| \le k - 1$*.*

2 Main results

For a graph G, let $V(G)$ and $E(G)$ denote the set of vertices and the set of edges of G, respectively. As usual, the *union* of two graphs G and H is the graph, denoted by $G \cup H$, with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H)$. Let mH be the disjoint union of m copies of a graph H. If M is a subset of edges of a graph G, the subgraph of G induced by M is denoted by $G[M]$. If M is the edge subset of G, then $G \setminus M$ denote the subgraph by deleting the edges of M. If $M = e$, we simply write $G \setminus e$ for $G \setminus \{e\}$. If $S \subseteq V(G)$, the subgraph of G induced by S is denoted by $G[S]$. For $S \subseteq V(G)$, we denote $G \setminus S$ the subgraph by deleting the vertices of S together with the edges incident with them from G. We denote by $E_G[X, Y]$ the set of edges of G with one end in X and the other end in Y. If $X = \{x\}$, we simply write $E[x, Y]$ for $E[\{x\}, Y]$. A subset M of $E(G)$ is called a *matching* of G if the edges of M satisfy that no two of them are adjacent in G. A matching M saturates a vertex v, or v is said to be *M-saturated*, if some edge of M is incident with v; otherwise, v is *M-unsaturated*. If every vertex of G is M-saturated, the matching M is *perfect*. M is a *maximum matching* if G has no matching M' with $|M'| > |M|$.

Lemma 5. *If G is a graph obtained from the complete graph K_n by deleting an edge set M and $\Delta(K_n[M]) \ge r$, then $\lambda_k(G) \le n - 1 - r$.*

Proof. Since $\Delta(K_n[M]) \ge r$, there exists at least one vertex, say v, such that $d_{K_n[M]}(v) \ge r$. Then $d_G(v) = n - 1 - d_{K_n[M]}(v) \le n - 1 - r$. So $\delta(G) \le d_G(v) \le n - 1 - r$. From Observation 1, $\lambda_k(G) \le \delta(G) \le n - 1 - r$. □

Corollary 1. *For every two integers n and k with $4 \le k \le n$, if k is even and M is an edge set of the complete graph K_n such that $\Delta(K_n[M]) \ge \frac{k}{2} + 1$, then $\kappa_k(K_n \setminus M) \le \lambda_k(K_n \setminus M) < n - \frac{k}{2} - 1$.*

Remark 1. From Corollary 1, if $\kappa_k(K_n \setminus M) = n - \frac{k}{2} - 1$ or $\lambda_k(K_n \setminus M) = n - \frac{k}{2} - 1$ for k even, then $\Delta(K_n[M]) \le \frac{k}{2}$.

In [15], we stated a useful Lemma for a general k.

Let $S = \{u_1, u_2, \cdots, u_k\} \subseteq V(G)$ and $\bar{S} = V(G) \setminus S = \{w_1, w_2, \cdots, w_{n-k}\}$. Denote by $\mathscr{T}$ a maximum set of edge-disjoint trees connecting S in G. Let $\mathscr{T}_1$ be the set of trees in $\mathscr{T}$ whose edges only belong to $E(G[S])$, and let $\mathscr{T}_2$ be the set of trees containing at least one edge of $E_G[S,\bar{S}]$. Thus, $\mathscr{T}=\mathscr{T}_1\cup\mathscr{T}_2$.

Lemma 6. *[15] Let G be a connected graph of order n, and $S\subseteq V(G)$ with $|S|=k$ $(3\leq k\leq n)$, and T be a Steiner tree connecting S. If $T\in\mathscr{T}_1$, then T uses exactly $k-1$ edges of $E(G[S])\cup E_G[S,\bar{S}]$; If $T\in\mathscr{T}_2$, then T uses at least k edges of $E(G[S])\cup E_G[S,\bar{S}]$.*

Lemma 7. *For every two integers n and k with $4\leq k\leq n$, if k is even and M is an edge set of the complete graph K_n such that $|M|\geq k$ and $\Delta(K_n[M])\geq 2$, then $\lambda_k(K_n\setminus M)<n-\frac{k}{2}-1$.*

Proof. Set $G=K_n\setminus M$. We claim that there is an $S\subseteq V(G)$ with $|S|=k$ such that $|M\cap\big(E(K_n[S])\cup E_{K_n}[S,\bar{S}]\big)|\geq k$ and $|M\cap\big(E(K_n[S])\big)|\geq 1$. Choose an edge set $M'\subseteq M$ such that $|M'|=k$. Suppose that $K_n[M']$ contains s independent edges and r connected components $C_1,\cdots,C_r$ such that $\Delta(C_i)\geq 2$ $(1\leq i\leq r)$. Set $|C_i|=n_i$ and $|E(C_i)|=m_i$. Then $m_i\geq n_i-1$. For each C_i $(1\leq i\leq r)$, choose one of the vertices with the maximum degree, say u_i. Set $X_i=V(C_i)\setminus u_i$.

If there exists some X_j such that $|E(K_n[X_j])|\geq 1$, then choose $X_i\subseteq S$ for all $1\leq i\leq r$. Since $|C_i|=n_i$ and $X_i=V(C_i)\setminus u_i$, we have $|X_i|=n_i-1$. By such a choosing, the number of the vertices belonging to S is $\sum_{i=1}^{r}|X_i|=\sum_{i=1}^{r}(n_i-1)\leq\sum_{i=1}^{r}m_i\leq k-s$. In addition, choose one endpoint of each independent edge into S. Till now, the total number of the vertices belonging to S is $\sum_{i=1}^{r}|X_i|+s\leq(k-s)+s=k$. Note that if $\sum_{i=1}^{r}|X_i|+s<k$, then we can add some other vertices in G into S such that $|S|=k$. Thus all edges of $E(C_i)$ and the s independent edges are put into $E(K_n[S])\cup E_{K_n}[S,\bar{S}]$, that is, all edges of M' belong to $E(K_n[S])\cup E_{K_n}[S,\bar{S}]$. So $|M\cap\big(E(K_n[S])\cup E_{K_n}[S,\bar{S}]\big)|\geq k$, as desired. Since $|E(K_n[X_j])|\geq 1$, it follows that $|M\cap\big(E(K_n[S])\big)|\geq 1$, as desired.

Suppose that $|E(K_n[X_i])|=0$ for all $1\leq i\leq r$. Then each C_i must be a star such that $|E(C_i)|\geq 2$. Recall that u_i is one of the vertices with the maximum degree in C_i. Pick up one vertex from $V(C_i)\setminus u_i$, say v_i. Put all the vertices of $Y_i=V(C_i)\setminus v_i$ into S, that is, $Y_i\subseteq S$. Thus $|Y_i|=n_i-1$. In addition, choose one endpoint of each independent edge into S. By such a choosing, the total number of the vertices belonging to S is $\sum_{i=1}^{r}|Y_i|+s=\sum_{i=1}^{r}(n_i-1)+s\leq\sum_{i=1}^{r}m_i+s\leq(k-s)+s=k$. Note that if $\sum_{i=1}^{r}|X_i|+s<k$ then we can add some other vertices in G into S such that $|S|=k$. Thus all edges of $E(C_i)$ and the s independent edge are put into $E(K_n[S])\cup E_{K_n}[S,\bar{S}]$, that is, and all edges of M' belong to $E(K_n[S])\cup E_{K_n}[S,\bar{S}]$. So $|M\cap\big(E(K_n[S])\cup E_{K_n}[S,\bar{S}]\big)|\geq k$, as desired. Since $|E(C_i)|\geq 2$, it follows that there is an edge $u_iw_i\in M\cap K_n[S]$ where $w_i\in V(C_i)\setminus v_i$, which implies that $|M\cap\big(E(K_n[S])\big)|\geq 1$, as desired.

From the above, we conclude that there exists an $S\subseteq V(G)$ with $|S|=k$ such that $|M\cap\big(E(K_n[S])\cup E_{K_n}[S,\bar{S}]\big)|\geq k$ and $|M\cap\big(E(K_n[S])\big)|\geq 1$. Since each tree $T\in\mathscr{T}_1$ uses $k-1$ edges in $E(G[S])\cup E_G[S,\bar{S}]$, it follows that $|\mathscr{T}_1|\leq(\binom{k}{2}-1)/(k-1)=$

$\frac{k}{2}-\frac{1}{k-1}$, which results in $|\mathscr{T}_1|\leq\frac{k}{2}-1$ since $|\mathscr{T}_1|$ is an integer. From Lemma 6, each tree $T\in\mathscr{T}_2$ uses at least k edges of $E(G[S])\cup E_G[S,\bar{S}]$. Thus $|\mathscr{T}_1|(k-1)+|\mathscr{T}_2|k\leq |E(G[S])|+|E_G[S,\bar{S}]|$, that is, $|\mathscr{T}_1|k+|\mathscr{T}_2|k\leq|\mathscr{T}_1|+\binom{k}{2}+k(n-k)-k$. So $\lambda_k(G)=|\mathscr{T}|=|\mathscr{T}_1|+|\mathscr{T}_2|\leq n-\frac{k}{2}-1-\frac{1}{k}<n-\frac{k}{2}-1$. ☐

Remark 2. From Lemmas 3 and 6, if $\kappa_k(K_n\backslash M)=n-\frac{k}{2}-1$ or $\lambda_k(K_n\backslash M)=n-\frac{k}{2}-1$ for k even and $2\leq\Delta(K_n[M])\leq\frac{k}{2}$, then $1\leq|M|\leq k-1$, where $M\subseteq E(K_n)$.

Lemma 8. *For every two integers n and k with $4\leq k\leq n$, if k is even and M is an edge set of the complete graph K_n such that $|M|\geq k$ and $\Delta(K_n[M])=1$, then $\kappa_k(K_n\setminus M)<n-\frac{k}{2}-1$.*

Proof. Since $\Delta(K_n[M])=1$, M is a matching in K_n. Since $|M|\geq k$, we can choose $M_1\subseteq M$ such that $|M_1|=k$. Let $M_1=\{u_iw_i|1\leq i\leq k\}$. Choose $S=\{u_1,u_2,\cdots,u_k\}$. We will show that $\kappa(S)<n-\frac{k}{2}-1$. Clearly, $|\bar{S}|=n-k$, and let $\bar{S}=\{w_1,w_2,\cdots,w_{n-k}\}$. Since each tree in $\mathscr{T}_2$ contains at least one vertex of $\bar{S}$, we have $|\mathscr{T}_2|\leq n-k$. By the definition of $\mathscr{T}_1$, we have $|\mathscr{T}_1|\leq\frac{k}{2}$. If $|\mathscr{T}_1|\leq\frac{k}{2}-2$, then $\lambda(S)=|\mathscr{T}|=|\mathscr{T}_1|+|\mathscr{T}_2|\leq(\frac{k}{2}-2)+(n-k)=n-\frac{k}{2}-2<n-\frac{k}{2}-1$, as desired. Let us assume $\frac{k}{2}-1\leq|\mathscr{T}_1|\leq\frac{k}{2}$.

Consider $|\mathscr{T}_1|=\frac{k}{2}-1$. Recall that $|\mathscr{T}_2|\leq n-k$. Furthermore, we claim that $|\mathscr{T}_2|\leq n-k-1$. Assume, to the contrary, that $|\mathscr{T}_2|=n-k$. Let $T_1,T_2,\cdots,T_{n-k}$ be the $n-k$ trees connecting S in $\mathscr{T}_2$. For each tree T_i $(1\leq i\leq n-k)$, this tree only occupy one vertex of $\bar{S}$, say w_i. Since $u_iw_i\in M_1$ $(1\leq i\leq k)$, namely, $u_iw_i\notin E(G)$, and each T_i $(1\leq i\leq k)$ is a tree connecting S in $\mathscr{T}_2$, it follows that this tree T_i must contain at least one edge in $G[S]=K_k$. So the trees $T_1,T_2,\cdots,T_k$ must use at least k edges in $G[S]$, and $|\mathscr{T}_1|=\frac{\binom{k}{2}-k}{k-1}=\frac{k-2}{2}-\frac{1}{k-1}$. Since $|\mathscr{T}_1|$ is an integer, we have $|\mathscr{T}_1|<\frac{k-2}{2}$, a contradiction. We conclude that $|\mathscr{T}_2|\leq n-k-1$, and hence $\lambda(S)=|\mathscr{T}|=|\mathscr{T}_1|+|\mathscr{T}_2|\leq(\frac{k}{2}-1)+(n-k-1)=n-\frac{k}{2}-2<n-\frac{k}{2}-1$, as desired.

Consider $|\mathscr{T}_1|=\frac{k}{2}$. We claim that $|\mathscr{T}_2|\leq n-k-2$. Assume, to the contrary, that $n-k-1\leq|\mathscr{T}_2|\leq n-k$. Since $|\mathscr{T}_1|=\frac{k}{2}$, it follows that each edge of $G[S]$ is occupied by some tree in $\mathscr{T}_1$, which implies that each tree in $\mathscr{T}_2$ only uses the edges of $E_G[S,\bar{S}]\cup E(G[\bar{S}])$. Suppose that T_1 is a tree in $\mathscr{T}_2$ connecting w_1. Since $u_1w_1\notin E(G)$, if T_1 contains three vertices of $\bar{S}$, then the remaining $n-k-3$ vertices in $\bar{S}$ must be contained in at most $n-k-3$ trees in $\mathscr{T}_2$, which results in $|\mathscr{T}_2|\leq(n-k-3)+1=n-k-2$, a contradiction. So we assume that the tree T_1 contains another vertex of $\bar{S}$ except w_1, say w_2. Recall that $k\geq4$. Then $|\bar{S}|\geq k\geq4$. By the same reason, there is another tree T_2 containing two vertices of $\bar{S}$, say w_3,w_4. Furthermore, the remaining $n-k-4$ vertices in $\bar{S}$ must be contained in at most $n-k-4$ trees in $\mathscr{T}_2$, which results in $|\mathscr{T}_2|\leq(n-k-4)+2=n-k-2$, a contradiction. We conclude that $|\mathscr{T}_2|\leq n-k-2$. Combining this with $|\mathscr{T}_1|=\frac{k}{2}$, we have $\lambda(S)=|\mathscr{T}|=|\mathscr{T}_1|+|\mathscr{T}_2|\leq\frac{k}{2}+(n-k-2)=n-\frac{k}{2}-2<n-\frac{k}{2}-1$, as desired. ☐

Lemma 9. *If n $(n \geq 4)$ is even and M is an edge set of the complete graph K_n such that $1 \leq |M| \leq n-1$ and $1 \leq \Delta(K_n[M]) \leq \frac{n}{2}$, then $G = K_n \setminus M$ contains $\frac{n-2}{2}$ edge-disjoint spanning trees.*

Proof. Let $\mathscr{P} = \bigcup_{i=1}^{p} V_i$ be a partition of $V(G)$ with $|V_i| = n_i$ $(1 \leq i \leq p)$, and $\mathcal{E}_p$ be the set of edges between distinct blocks of $\mathscr{P}$ in G. It suffices to show that $|\mathcal{E}_p| \geq \frac{n-2}{2}(|\mathscr{P}|-1)$ so that we can use Theorem 1.

The case $p = 1$ is trivial by Theorem 1, thus we assume $p \geq 2$. For $p = 2$, we have $\mathscr{P} = V_1 \cup V_2$. Set $|V_1| = n_1$. Clearly, $|V_2| = n - n_1$. Since $\Delta(K_n[M]) \leq \frac{n}{2}$, it follows that $\delta(G) = n - 1 - \Delta(K_n[M]) \geq n - 1 - \frac{n}{2} = \frac{n-2}{2}$. Therefore, if $n_1 = 1$ then $|\mathcal{E}_2| = |E_G[V_1, V_2]| \geq \frac{n-2}{2} = \frac{n-2}{2}$. Suppose $n_1 \geq 2$. Then $|\mathcal{E}_2| = |E_G[V_1, V_2]| \geq \binom{n}{2} - (n-1) - \binom{n_1}{2} - \binom{n-n_1}{2} = -n_1^2 + nn_1 - n + 1$. Since $2 \leq n_1 \leq n-2$, one can see that $|\mathcal{E}_2|$ obtain its minimum value when $n_1 = 2$ or $n_1 = n-2$. Thus $|\mathcal{E}_2| \geq n - 3 \geq \frac{n-2}{2}$ since $n \geq 4$. The result follows from Theorem 1.

Consider the remaining case p with $3 \leq p \leq n$. Since $|\mathcal{E}_p| \geq \binom{n}{2} - |M| - \sum_{i=1}^{p}\binom{n_i}{2} \geq \binom{n}{2} - (n-1) - \sum_{i=1}^{p}\binom{n_i}{2} = \binom{n-1}{2} - \sum_{i=1}^{p}\binom{n_i}{2}$, we only need to show $\binom{n-1}{2} - \sum_{i=1}^{p}\binom{n_i}{2} \geq \frac{n-2}{2}(p-1)$, that is, $(n-p)\frac{n-2}{2} \geq \sum_{i=1}^{p}\binom{n_i}{2}$. Let $f(n_1, n_2, \cdots, n_p) = \sum_{i=1}^{p}\binom{n_i}{2}$. Then we only need to prove that $(n-p)\frac{n-2}{2} \geq f_{max}(n_1, n_2, \cdots, n_p)$. Because $f(n_1, n_2, \cdots, n_p) = \sum_{i=1}^{p}\binom{n_i}{2}$ achieves its maximum value when $n_1 = n_2 = \cdots = n_{p-1} = 1$ and $n_p = n - p + 1$, we need inequality $(n-p)\frac{n-2}{2} \geq \binom{1}{2}(p-1) + \binom{n-p+1}{2}$, namely, $(n-p)\frac{p-3}{2} \geq 0$. It is easy to see that the inequality holds since $3 \leq p \leq n$. Thus, $|\mathcal{E}_p| \geq \binom{n}{2} - |M| - \sum_{i=1}^{p}\binom{n_i}{2} \geq \frac{n-2}{2}(p-1)$. From Theorem 1, there exist $\frac{n-2}{2}$ edge-disjoint spanning trees in G, as desired. ☐

Lemma 10. *Let k, n be two integers with $4 \leq k \leq n$, and M is an edge set of the complete graph K_n satisfying $\Delta(K_n[M]) = 1$. Then*

(1) If $|M| = k-1$, then $\kappa_k(K_n \setminus M) \geq n - \frac{k}{2} - 1$;

(2) If $|M| = \lfloor \frac{n}{2} \rfloor$, then $\lambda_k(K_n \setminus M) \geq n - \frac{k}{2} - 1$.

Proof. (1) Set $G = K_n \setminus M$. Since $\Delta(K_n[M]) = 1$, it follows that M is a matching of K_n. By the definition of $\kappa_k(G)$, we need to show that $\kappa(S) \geq n - \frac{k}{2} - 1$ for any $S \subseteq V(G)$.

Case 1. There exists no u, w in S such that $uw \in M$.

Without loss of generality, let $S = \{u_1, u_2, \cdots, u_k\}$ such that $u_1, u_2, \cdots, u_r$ are M-saturated but $u_{r+1}, u_{r+2}, \cdots, u_k$ are M-unsaturated. Let $M_1 = \{u_iw_i | 1 \leq i \leq r\} \subseteq M$. Since $|M| = k-1$, it follows that $0 \leq r \leq k-1$. In this case, $u_iu_j \notin M$ $(1 \leq i, j \leq r)$. Clearly, $G[S]$ is a clique of order k, and choose a path $P = u_1u_2\cdots u_ru_{r+1}$ in $G[S]$. Let $G' = G \setminus E(P)$. Then $G'[S] = K_k \setminus E(P)$. Since $|E(P)| = r \leq k-1$ and $\Delta(K_k[E(P)]) = 2 \leq \frac{k}{2}$, it follows that $G'[S]$ contains $\frac{k-2}{2}$ spanning trees, which are also $\frac{k-2}{2}$ internally disjoint connecting S. These trees together with $T_i = u_1w_i \cup u_2w_i \cup u_{i-1}w_i \cup u_{i+1}w_i \cup \cdots \cup u_kw_i \cup u_iu_{i+1}$ $(1 \leq i \leq r)$ (see Figure 1 (a)) and $T_j = u_1v_j \cup u_2v_j \cup \cdots \cup u_kv_j$ where $v_j = \bar{S} \setminus \{w_1, w_2, \cdots, w_r\} = \{v_1, v_2, \cdots, v_{n-k-r}\}$

form $\frac{k-2}{2}+r+(n-k-r)=n-\frac{k}{2}-1$ internally disjoint trees connecting S, namely, $\kappa(S)\geq n-\frac{k}{2}-1$, as desired.

Case 2. There exist u, w in S such that $uw\in M$.

Without loss of generality, let $S=\{u_1,u_2,\cdots,u_r,u_{r+1},u_{r+2},\cdots,u_{r+s},u_{r+s+1},\cdots,u_{k-r},w_1,w_2,\cdots,w_r\}$ such that the vertices $u_1,u_2,\cdots,u_{r+s},w_1,w_2,\cdots,w_r$ are all M-saturated and $u_iw_i\in M$ $(1\leq i\leq r)$. Set $M_1=\{u_iw_i|1\leq i\leq r\}$. In this case, $r\geq 1$ and $2r+s\leq k$. Since $|M|=k-1$, it follows that $r+s\leq k-1$ and $s\leq k-2$.

First, we consider $2r+s=k$. Since k is even, it follows that s is even. If $s=0$, then $r=\frac{k}{2}$. Thus $S=\{u_1,u_2,\cdots,u_{\frac{k}{2}},w_1,w_2,\cdots,w_{\frac{k}{2}}\}$. Clearly, $M_1=\{u_iw_i|1\leq i\leq \frac{k}{2}\}$, $|M_1|=\frac{k}{2}\leq k-1$ and $\Delta(K_n[M_1])=1<\frac{k}{2}$. By Lemma 9, $G[S]$ contains $\frac{k-2}{2}$ spanning trees, which are also $\frac{k-2}{2}$ internally disjoint trees connecting S. These trees together with $T_j=u_1v_j\cup u_2v_j\cup\cdots\cup u_{\frac{k}{2}}v_j\cup w_1v_j\cup w_2v_j\cup\cdots\cup w_{\frac{k}{2}}v_j$ form $\frac{k-2}{2}+(n-k)$ internally disjoint trees, where $v_j\in\bar{S}=\{v_1,v_2,\cdots,v_{n-k}\}$, namely, $\kappa(S)\geq n-\frac{k}{2}-1$.



(a) (b) (c) (d)

Figure 1. Graphs for (1) of Lemma 10.

Consider $s=2$. Since $2r+s=k$, we have $r=\frac{k-2}{2}$. If $k=4$, then $r=1$ and hence $S=\{u_1,u_2,u_3,w_1\}$. Clearly, $M_1=\{u_1w_1\}$, and the trees $T_1=u_1u_2\cup u_1w_2\cup w_1w_2\cup u_3w_2$ and $T_2=u_1u_3\cup u_2u_3\cup u_2w_1$ and $T_3=u_1w_3\cup u_2w_3\cup w_1w_3\cup u_3w_1$ are three internally disjoint spanning trees connecting S; see Figure 1 (c). These trees together with $T_j=u_1v_j\cup u_2v_j\cup u_3v_j\cup w_1v_j$ form $3+(n-6)$ internally disjoint trees, where $v_j\in\bar{S}\setminus\{w_2,w_3\}=\{v_1,v_2,\cdots,v_{n-6}\}$, namely, $\kappa(S)\geq n-3=n-\frac{k}{2}-1$. Assume $k\geq 6$. Then $r\geq 2$, $S=\{u_1,u_2,\cdots,u_{\frac{k+2}{2}},w_1,w_2,\cdots,w_{\frac{k-2}{2}}\}$ and $M_1=\{u_iw_i|1\leq i\leq\frac{k-2}{2}\}$. Clearly, the trees $T_1=u_1w_{\frac{k}{2}}\cup u_2w_{\frac{k}{2}}\cup\cdots\cup u_{\frac{k-2}{2}}w_{\frac{k}{2}}\cup$

$u_{\frac{k+2}{2}}w_{\frac{k}{2}} \cup u_2u_{\frac{k}{2}} \cup w_1w_{\frac{k}{2}} \cup w_2w_{\frac{k}{2}} \cup \cdots \cup w_{\frac{k-2}{2}}w_{\frac{k}{2}}$ and $T_2 = u_1w_{\frac{k+2}{2}} \cup u_2w_{\frac{k+2}{2}} \cup \cdots \cup u_{\frac{k}{2}}w_{\frac{k+2}{2}} \cup u_1u_{\frac{k+2}{2}} \cup w_1w_{\frac{k+2}{2}} \cup w_2w_{\frac{k+2}{2}} \cup \cdots \cup w_{\frac{k-2}{2}}w_{\frac{k+2}{2}}$ are two internally disjoint trees connecting S; see Figure 1 (d). Let $M_2 = M_1 \cup \{u_1u_{\frac{k+2}{2}}, u_2u_{\frac{k}{2}}\}$. Then $|M_2| = |M_1| + 2 = \frac{k-2}{2} + 2 = \frac{k+2}{2} < k-1$ and $\Delta(K_n[M_2]) = 2 \leq \frac{k}{2}$, which implies that $G[S] \setminus \{u_1u_{\frac{k+2}{2}}, u_2u_{\frac{k}{2}}\} = K_k \setminus M_2$ contains $\frac{k-2}{2}$ spanning trees by Lemma 9, which are also $\frac{k-2}{2}$ internally disjoint trees connecting S. These trees together with T_1, T_2 and the trees $T_j = u_1v_j \cup u_2v_j \cup \cdots \cup u_{\frac{k+2}{2}}v_j \cup w_1v_j \cup w_2v_j \cup \cdots \cup u_{\frac{k-2}{2}}v_j$ are $\frac{k-2}{2}+2+(n-k-2)$ internally disjoint trees connecting S, where $v_j \in \bar{S}\setminus\{w_{\frac{k}{2}}, w_{\frac{k+2}{2}}\} = \{v_1, v_2, \cdots, v_{n-k-2}\}$, namely, $\kappa(S) \geq n - \frac{k}{2} - 1$.

Consider the remaining case s with $4 \leq s \leq k-2$. Clearly, there exists a cycle of order s containing $u_{r+1}, u_{r+2}, \cdots, u_{r+s}$ in $K_k \setminus M_1$, say $C_s = u_{r+1}u_{r+2}\cdots u_{r+s}u_{r+1}$. Set $M' = M_1 \cup E(C_s)$. Then $|M'| = r + s \leq k-1$ and $\Delta(K_n[M']) = 2 \leq \frac{k}{2}$, which implies that $G \setminus E(C_s) = K_k \setminus M'$ contains $\frac{k-2}{2}$ spanning trees by Lemma 9. These trees together with the trees $T_{r+j} = u_1w_{r+j} \cup u_2w_{r+j} \cup \cdots \cup u_{r+j-1}w_{r+j} \cup u_{r+j+1}w_{r+j} \cup \cdots \cup u_{r+s}w_{r+j} \cup u_{r+j}u_{r+j+1} \cup w_1w_{r+j} \cup w_2w_{r+j} \cup \cdots \cup w_rw_{r+j}$ $(1 \leq j \leq s)$ form $\frac{k-2}{2} + s$ internally disjoint trees; see Figure 2 (b) (note that $u_{r+s} = u_{k-r}$). These trees together with $T'_j = u_1v_j \cup u_2v_j \cup \cdots \cup u_{r+s}v_j \cup w_1v_j \cup \cdots \cup w_rv_j$ form $\frac{k-2}{2} + s + (n-2r-2s) = n - \frac{k}{2} - 1$ internally disjoint trees connecting S where $v_j \in \bar{S} \setminus \{w_{r+1}, w_{r+2}, \cdots, w_{r+s}\} = \{v_1, v_2, \cdots, v_{n-2r-2s}\}$, namely, $\kappa(S) \geq n - \frac{k}{2} - 1$, as desired.

Next, assume $2r + s < k$. Then $S = \{u_1, u_2, \cdots, u_{r+s}, u_{r+s+1}, \cdots, u_{k-r}, w_1, w_2, \cdots, w_r\}$ and $r+s+1 \leq k-r$. If $s = 0$, then $S = \{u_1, u_2, \cdots, u_{k-r}, w_1, w_2, \cdots, w_r\}$. Clearly, $M_1 = \{u_iw_i | 1 \leq i \leq r\}$, $|M_1| = r \leq k-1$ and $\Delta(K_n[M_1]) = 1 < \frac{k}{2}$. By Lemma 9, $G[S]$ contains $\frac{k-2}{2}$ spanning trees. These trees together with $T_j = u_1v_j \cup u_2v_j \cup \cdots \cup u_{n-r}v_j \cup w_1v_j \cup w_2v_j \cup \cdots \cup w_rv_j$ form $\frac{k-2}{2} + (n-k)$ internally disjoint trees, where $v_j \in \bar{S} = \{v_1, v_2, \cdots, v_{n-k}\}$, namely, $\kappa(S) \geq n - \frac{k}{2} - 1$. Assume $s \geq 1$. Clearly, there exists a path of length s containing $u_{r+1}, u_{r+2}, \cdots, u_{r+s}, u_{r+s+1}$ in $G[S]$, say $P_s = u_{r+1}u_{r+2}\cdots u_{r+s}u_{r+s+1}$. Set $M' = M_1 \cup E(P_s)$. Then $|M'| = r + s \leq k-1$ and $\Delta(K_n[M']) = 2 \leq \frac{k}{2}$, which implies that $G[S] \setminus E(P_s) = K_k \setminus M'$ contains $\frac{k-2}{2}$ spanning trees by Lemma 9, which are also $\frac{k-2}{2}$ internally disjoint trees connecting S. These trees together with the trees $T_{r+j} = u_1w_{r+j} \cup u_2w_{r+j} \cup \cdots \cup u_{r+j-1}w_{r+j} \cup u_{r+j+1}w_{r+j} \cup \cdots \cup u_{k-r}w_{r+j} \cup u_{r+j}u_{r+j+1} \cup w_1w_{r+j} \cup w_2w_{r+j} \cup \cdots \cup w_rw_{r+j}$ $(1 \leq j \leq s)$ form $\frac{k-2}{2} + s$ internally disjoint trees; see Figure 1 (b). These trees together with $T'_j = u_1v_j \cup u_2v_j \cup \cdots \cup u_{k-r}v_j \cup w_1v_j \cup w_2v_j \cup \cdots \cup w_rv_j$ form $\frac{k-2}{2} + s + (n-k+r) - (r+s) = n - \frac{k}{2} - 1$ internally disjoint trees connecting S where $v_j \in \bar{S} \setminus \{w_{r+1}, w_{r+2}, \cdots, w_{r+s}\} = \{v_1, v_2, \cdots, v_{n-k-s}\}$, namely, $\kappa(S) \geq n - \frac{k}{2} - 1$, as desired.

We conclude that $\kappa(S) \geq n - \frac{k}{2} - 1$ for any $S \subseteq V(G)$. From the arbitrariness of S, it follows that $\kappa_k(G) \geq n - \frac{k}{2} - 1$.

(2) Set $G = K_n \setminus M$. Assume that n is even. Thus M is a perfect matching of

K_n, and all vertices of G are M-saturated. By the definition of $\lambda_k(G)$, we need to show that $\lambda(S) \geq n - \frac{k}{2} - 1$ for any $S \subseteq V(G)$.

Case 3. There exists no u, w in S such that $uw \in M$.

Without loss of generality, let $S = \{u_1, u_2, \cdots, u_k\}$. In this case, $u_iu_j \notin M$ $(1 \leq i, j \leq k)$. Let $M_1 = \{u_iw_i | 1 \leq i \leq k\} \subseteq M = \{u_iw_i | 1 \leq i \leq \frac{n}{2}\}$. Clearly, $w_i \notin S$ $(1 \leq i \leq \frac{n}{2})$ and $u_j \notin S$ $(k+1 \leq j \leq \frac{n}{2})$. Since $G[S]$ is a clique of order k, it follows that there are $\frac{k}{2}$ edge-disjoint spanning trees in $G[S]$, which are also $\frac{k}{2}$ edge-disjoint trees connecting S. These trees together with $T_i = u_1w_i \cup u_2w_i \cup u_{i-1}w_i \cup u_{i+1}w_i \cup \cdots \cup u_kw_i \cup u_iw_k \cup w_iw_k$ $(1 \leq i \leq k-1)$ (see Figure 2 (a)) and $T_j' = u_1u_j \cup u_2u_j \cup \cdots \cup u_ku_j$ $(k+1 \leq j \leq \frac{n}{2})$ and $T_j'' = u_1w_j \cup u_2w_j \cup \cdots \cup u_kw_j$ $(k+1 \leq j \leq \frac{n}{2})$ form $\frac{k}{2} + (k-1) + (n-2k) = n - \frac{k}{2} - 1$ edge-disjoint trees connecting S, namely, $\lambda(S) \geq n - \frac{k}{2} - 1$, as desired.

Case 4. There exist u, w in S such that $uw \in M$.

Without loss of generality, let $S = \{u_1, u_2, \cdots, u_{r+s}, w_1, w_2, \cdots, w_r\}$ with $|S| = k = 2r + s$, where $1 \leq r \leq \frac{k}{2}$ and $0 \leq s \leq k-2$. Set $M_1 = \{u_iw_i | 1 \leq i \leq r\} \subseteq M = \{u_iw_i | 1 \leq i \leq \frac{n}{2}\}$. We claim that $r + s \leq k - 1$. Otherwise, let $r + s = k$. Combining this with $2r + s = k$, we have $r = 0$, a contradiction. Since $k = 2r + s$ and k is even, it follows that s is even.



(a) (b)

Figure 2. Graphs for (2) of Lemma 10.

If $s = 0$, then $r = \frac{k}{2}$. Clearly, $S = \{u_1, u_2, \cdots, u_{\frac{k}{2}}, w_1, w_2, \cdots, w_{\frac{k}{2}}\}$ and $M_1 = M = \{u_iw_i | 1 \leq i \leq \frac{k}{2}\}$. In addition, $|M_1| \leq \frac{k}{2} < k - 1$ and $\Delta(M \cap K_n[S]) = 1 < \frac{k}{2}$. Then $G[S]$ contains $\frac{k-2}{2}$ edge-disjoint spanning trees connecting S by Lemma 9. These trees together with $T_i = u_1u_i \cup u_2u_i \cup \cdots \cup u_{\frac{k}{2}}u_i \cup w_1u_i \cup w_2u_i \cup \cdots \cup w_{\frac{k}{2}}u_i$ $(\frac{k}{2}+1 \leq i \leq \frac{n}{2})$ and $T_i' = u_1w_i \cup u_2w_i \cup \cdots \cup u_{\frac{k}{2}}w_i \cup w_1w_i \cup w_2w_i \cup \cdots \cup w_{\frac{k}{2}}w_i$ $(\frac{k}{2}+1 \leq i \leq \frac{n}{2})$ form $n - \frac{k}{2} - 1$ edge-disjoint trees connecting S, namely, $\lambda(S) \geq n - \frac{k}{2} - 1$.

If $s = 2$, then $r = \frac{k-2}{2}$. Then $S = \{u_1, u_2, \cdots, u_{\frac{k+2}{2}}, w_1, w_2, \cdots, w_{\frac{k-2}{2}}\}$ and $M_1 = \{u_iw_i | 1 \leq i \leq \frac{k-2}{2}\} \subseteq M$. If $k = 4$, then $r = 1$ and hence $S = \{u_1, u_2, u_3, w_1\}$. Clearly, $M_1 = \{u_1w_1\}$, and the trees $T_1 = u_1u_2 \cup u_1w_2 \cup w_1w_2 \cup u_3w_2$ and $T_2 = u_1u_3 \cup u_2u_3 \cup u_2w_1$ and $T_3 = u_1w_3 \cup u_2w_3 \cup w_1w_3 \cup u_3w_1$ are three edge-disjoint spanning trees connecting S; see Figure 1 (c). These trees together with $T_j = u_1u_j \cup u_2u_j \cup u_3u_j \cup w_1u_j$ $(4 \leq k \leq \frac{n}{2})$ and $T_j' = u_1w_j \cup u_2w_j \cup u_3w_j \cup w_1u_j$ $(4 \leq k \leq \frac{n}{2})$

form $3+(n-6)$ edge-disjoint trees connecting S, namely, $\lambda(S)\geq n-3=n-\frac{k}{2}-1$, as desired. Assume $k\geq 6$. Then $r\geq 2$, $S=\{u_1,u_2,\cdots,u_{\frac{k+2}{2}},w_1,w_2,\cdots,w_{\frac{k-2}{2}}\}$ and $M_1=\{u_iw_i|1\leq i\leq \frac{k-2}{2}\}$. Clearly, the trees $T_1=u_1w_{\frac{k}{2}}\cup u_2w_{\frac{k}{2}}\cup\cdots\cup u_{\frac{k-2}{2}}w_{\frac{k}{2}}\cup u_{\frac{k+2}{2}}w_{\frac{k}{2}}\cup u_2u_{\frac{k}{2}}\cup w_1w_{\frac{k}{2}}\cup w_2w_{\frac{k}{2}}\cup\cdots\cup w_{\frac{k-2}{2}}w_{\frac{k}{2}}$ and $T_2=u_1w_{\frac{k+2}{2}}\cup u_2w_{\frac{k+2}{2}}\cup\cdots\cup u_{\frac{k}{2}}w_{\frac{k+2}{2}}\cup u_1u_{\frac{k+2}{2}}\cup w_1w_{\frac{k+2}{2}}\cup w_2w_{\frac{k+2}{2}}\cup\cdots\cup w_{\frac{k-2}{2}}w_{\frac{k+2}{2}}$ are two edge-disjoint trees connecting S; see Figure 1 (d). Let $M_2=M_1\cup\{u_1u_{\frac{k+2}{2}},u_2u_{\frac{k}{2}}\}$. Then $|M_2|=|M_1|+2=\frac{k-2}{2}+2=\frac{k+2}{2}<k-1$ and $\Delta(K_n[M_2])=2\leq\frac{k}{2}$, which implies that $G[S]\setminus\{u_1u_{\frac{k+2}{2}},u_2u_{\frac{k}{2}}\}=K_k\setminus M_2$ contains $\frac{k-2}{2}$ spanning trees by Lemma 9. These trees together with T_1,T_2 and the trees $T_j=u_1u_j\cup u_2u_j\cup\cdots\cup u_{\frac{k+2}{2}}u_j\cup w_1u_j\cup w_2u_j\cup\cdots\cup u_{\frac{k-2}{2}}u_j$ ($\frac{k}{2}+2\leq j\leq\frac{n}{2}$) and $T_j=u_1w_j\cup u_2w_j\cup\cdots\cup u_{\frac{k+2}{2}}w_j\cup w_1w_j\cup w_2w_j\cup\cdots\cup u_{\frac{k-2}{2}}w_j$ ($\frac{k}{2}+2\leq j\leq\frac{n}{2}$) are $\frac{k-2}{2}+2+(n-k-2)$ edge-disjoint trees connecting S, namely, $\lambda(S)\geq n-\frac{k}{2}-1$, as desired.

Consider the remaining case s with $4\leq s\leq k-2$. Clearly, there exists a cycle of order s containing $u_{r+1},u_{r+2},\cdots,u_{r+s}$ in $K_k\setminus M_1$, say $C_s=u_{r+1}u_{r+2}\cdots u_{r+s}u_{r+1}$. Set $M'=M_1\cup E(C_s)$. Then $|M'|=r+s\leq k-1$ and $\Delta(K_n[M'])=2\leq\frac{k}{2}$, which implies that $G\setminus E(C_s)$ contains $\frac{k-2}{2}$ edge-disjoint spanning trees connecting S by Lemma 9. These trees together with the trees $T_{r+j}=u_1w_{r+j}\cup u_2w_{r+j}\cup\cdots\cup u_{r+j-1}w_{r+j}\cup u_{r+j+1}w_{r+j}\cup\cdots\cup u_{r+s}w_{r+j}\cup u_{r+j}u_{r+j+1}\cup w_1w_{r+j}\cup w_2w_{r+j}\cup\cdots\cup w_rw_{r+j}$ ($1\leq j\leq s$) form $\frac{k-2}{2}+s$ edge-disjoint trees; see Figure 2 (b). These trees together with $T_i'=u_1u_i\cup u_2u_i\cup\cdots\cup u_{r+s}u_i\cup w_1u_i\cup\cdots\cup w_ru_i$ ($r+s+1\leq i\leq\frac{n}{2}$) and $T_i''=u_1w_i\cup u_2w_i\cup\cdots\cup u_{r+s}w_i\cup w_1w_i\cup\cdots\cup w_rw_i$ ($r+s+1\leq i\leq\frac{n}{2}$) form $(n-2r-2s)+(\frac{k-2}{2}+s)=n-\frac{k}{2}-1$ edge-disjoint trees connecting S since $2r+s=k$, namely, $\lambda(S)\geq n-\frac{k}{2}-1$, as desired.

We conclude that $\lambda(S)\geq n-\frac{k}{2}-1$ for any $S\subseteq V(G)$. From the arbitrariness of S, it follows that $\lambda_k(G)\geq n-\frac{k}{2}-1$ For n odd, M is a maximum matching and we can also check that $\lambda_k(G)\geq n-\frac{k}{2}-1$ similarly. ☐

Lemma 11. *Let n and k be two integers such that k is even and $4\leq k\leq n$. If M is an edge set of the complete graph K_n such that $|M|=k-1$, and $2\leq\Delta(K_n[M])\leq\frac{k}{2}$, then $\kappa_k(K_n\setminus M)\geq n-\frac{k}{2}-1$.*

Proof. For $n=k$, there are $\frac{n-2}{2}$ edge-disjoint spanning trees by Lemma 9, namely, $\lambda_n(G)\geq\frac{n-2}{2}$. So from now on, we assume $n\geq k+1$. Let $S=\{u_1,u_2,\cdots,u_k\}\subseteq V(G)$ and $\bar{S}=V(G)\setminus S=\{w_1,w_2,\cdots,w_{n-k}\}$. We have the following two cases to consider.

Case 1. $M\subseteq E(K_n[S])\cup E(K_n[\bar{S}])$.

Let $M'=M\cap E(K_n[S])$ and $M''=M\cap E(K_n[\bar{S}])$. Then $|M'|+|M''|=|M|=k-1$ and $0\leq|M'|,|M''|\leq k-1$. We can regard $G[S]$ as a complete graph K_k by deleting $|M'|$ edges. Since $2\leq\Delta(K_n[M])\leq\frac{k}{2}$ and $M'\subseteq M$, it follow that $\Delta(K_n[M'])\leq\Delta(K_n[M])\leq\frac{k}{2}$. From Lemma 9, there exist $\frac{k-2}{2}$ edge-disjoint spanning trees in $G[S]$. Actually, these $\frac{k-2}{2}$ edge-disjoint spanning trees are all internally disjoint trees connecting S in $G[S]$. All these trees together with the trees $T_i = w_iu_1 \cup w_iu_2 \cup \cdots \cup w_iu_k$ $(1 \leq i \leq n-k)$ form $\frac{k-2}{2} + (n-k) = n - \frac{k}{2} - 1$ internally disjoint trees connecting S, namely, $\kappa(S) \geq n - \frac{k}{2} - 1$. From the arbitrariness of S, we have $\kappa_k(G) \geq n - \frac{k}{2} - 1$, as desired.

Case 2. $M \nsubseteq E(K_n[S]) \cup E(K_n[\bar{S}])$.

In this case, there exist some edges of M in $E_{K_n}[S, \bar{S}]$. Let $M' = M \cap E(K_n[S])$, $M'' = M \cap E(K_n[\bar{S}])$, and $|M'| = m_1$ and $|M''| = m_2$. Clearly, $0 \leq m_i \leq k-2$ $(i = 1, 2)$. For $w_i \in \bar{S}$, let $|E_{K_n[M]}[w_i, S]| = x_i$, where $1 \leq i \leq n-k$. Without loss of generality, let $x_1 \geq x_2 \geq \cdots \geq x_{n-k}$. Because there exist some edges of M in $E_{K_n}[S, \bar{S}]$, we have $x_1 \geq 1$. Since $2 \leq \Delta(K_n[M]) \leq \frac{k}{2}$, $x_i = |E_{K_n[M]}[w_i, S]| \leq d_{K_n[M]}(w_i) \leq \Delta(K_n[M]) \leq \frac{k}{2}$ for $1 \leq i \leq n-k$. We claim that there exists at most one vertex in $K_n[M]$ such that its degree is $\frac{k}{2}$. Assume, to the contrary, that there are two vertices, say w and w', such that $d_{K_n[M]}(w) = d_{K_n[M]}(w') = \frac{k}{2}$. Then $|M| \geq d_{K_n[M]}(w) + d_{K_n[M]}(w') \geq \frac{k}{2} + \frac{k}{2} = k$, contradicting to $|M| = k-1$. We conclude that there exists at most one vertex in $K_n[M]$ such that its degree is $\frac{k}{2}$. Recall that $x_{n-k} \leq x_{n-k-1} \leq \cdots \leq x_2 \leq x_1 \leq \frac{k}{2}$. So $x_1 = \frac{k}{2}$ and $x_i \leq \frac{k-2}{2}$ $(2 \leq i \leq n-k)$, or $x_i \leq \frac{k-2}{2}$ $(1 \leq i \leq n-k)$. Since $|E_{K_n[M]}[w_i, S]| = x_i$, we have $|E_G[w_i, S]| = k - x_i$. Since $2 \leq \Delta(K_n[M]) \leq \frac{k}{2}$, it follows that $\delta(G[S]) \geq k - 1 - \frac{k}{2} = \frac{k-2}{2}$.

Our basic idea is to seek for some edges in $G[S]$, and combine them with the edges of $E_G[S, \bar{S}]$ to form $n-k$ internally disjoint trees, say $T_1, T_2, \cdots, T_{n-k}$, with their roots $w_1, w_2, \cdots, w_{n-k}$, respectively. Let $G' = G \setminus \bigcup_{j=1}^{n-k} E(T_j)$. We will prove that $G'[S]$ satisfies the conditions of Lemma 9 so that $G'[S]$ contains $\frac{k-2}{2}$ edge-disjoint trees, which are also $\frac{k-2}{2}$ internally disjoint trees connecting S. These trees together with $T_1, T_2, \cdots, T_{n-k}$ are $n - \frac{k}{2} - 1$ our desired trees, namely, $\kappa(S) \geq n - \frac{k}{2} - 1$. So we can complete our proof by the arbitrariness of S.

For $w_1 \in \bar{S}$, without loss of generality, let $S = S_1^1 \cup S_2^1$ and $S_1^1 = \{u_1, u_2, \cdots, u_{x_1}\}$ such that $u_jw_1 \in M$ for $1 \leq j \leq x_1$. Set $S_2^1 = S \setminus S_1^1 = \{u_{x_1+1}, u_{x_1+2}, \cdots, u_k\}$. Then $u_jw_1 \in E(G)$ for $x_1+1 \leq j \leq k$. One can see that the tree $T_1' = w_1u_{x_1+1} \cup w_1u_{x_1+2} \cup \cdots \cup w_1u_k$ is a Steiner tree connecting S_2^1. Our current idea is to seek for x_1 edges in $E_G[S_1^1, S_2^1]$ and add them to T_1' to form a Steiner tree connecting S. For each $u_j \in S_1^1$ $(1 \leq j \leq x_1)$, we claim that $|E_G[u_j, S_2^1]| \geq 1$. Otherwise, let $|E_G[u_j, S_2^1]| = 0$. Then $|E_{K_n[M]}[u_j, S_2^1]| = k - x_1$ and hence $|M| \geq |E_{K_n[M]}[u_j, S_2^1]| + d_{K_n[M]}(w_1) \geq (k - x_1) + x_1 = k$, which contradicts to $|M| = k-1$. We conclude that for each $u_j \in S_1^1$ $(1 \leq j \leq x_1)$ there is at least one edge in G connecting it to a vertex of S_2^1. Choose the one with the smallest subscript among all the vertices of S_1^1 with the maximum degree in $G[S]$, say u_1'. Then we search for the vertex adjacent to u_1' with the smallest subscript among all the vertices of S_2^1 with the maximum degree in $G[S]$, say u_1''. Let $e_{11} = u_1'u_1''$. Consider the graph $G_{11} = G \setminus e_{11}$, and pick up the one with the smallest subscript among all the vertices of $S_1^1 \setminus u_1'$ with the maximum degree in $G_{11}[S]$, say u_2'. Then we search for the vertex adjacent to u_2' with the smallest subscript among all the vertices of S_2^1 with the maximum degree in $G_{11}[S]$, say u_2''. Set $e_{12} = u_2'u_2''$.

Consider the graph $G_{12} = G_{11} \setminus e_{12} = G \setminus \{e_{11}, e_{12}\}$. Choose the one with the smallest subscript among all the vertices of $S_1^1 \setminus \{u_1', u_2'\}$ with the maximum degree in $G_{12}[S]$, say u_3', and search for the vertex adjacent to u_3' with the smallest subscript among all the vertices of S_2^1 with the maximum degree in $G_{12}[S]$, say u_3''. Put $e_{13} = u_3'u_3''$. Consider the graph $G_{13} = G_{12} \setminus e_{11} = G \setminus \{e_{11}, e_{12}, e_{13}\}$. For each $u_j \in S_1^1$ $(1 \leq j \leq x_1)$, we proceed to find $e_{14}, e_{15}, \cdots, e_{1x_1}$ in the same way, and obtain graphs $G_{1j} = G \setminus \{e_{11}, e_{12}, \cdots, e_{1(j-1)}\}$ $(1 \leq j \leq x_1)$. Let $M_1 = \{e_{11}, e_{12}, \cdots, e_{1x_1}\}$ and $G_1 = G \setminus M_1$. Thus the tree $T_1 = w_1u_{x_1+1} \cup w_1u_{x_1+2} \cup \cdots \cup w_1u_k \cup e_{11} \cup e_{12} \cup \cdots \cup e_{1x_1}$ is our desired tree.

Let us now prove the following claim.

Claim 1. $\delta(G_1[S]) \geq \frac{k-2}{2}$.

Proof of Claim 1. Assume, to the contrary, that $\delta(G_1[S]) \leq \frac{k-4}{2}$. Then there exists a vertex $u_p \in S$ such that $d_{G_1[S]}(u_p) \leq \frac{k-4}{2}$. If $u_p \in S_2^1$, then by our procedure $d_{G[S]}(u_p) = d_{G_1[S]}(u_p) + 1 \leq \frac{k-2}{2}$, which implies that $d_{M \cap K_n[S]}(u_p) \geq k - 1 - \frac{k-2}{2} = \frac{k}{2}$. Since $w_1u_p \in M$, it follows that $d_{K_n[M]}(u_p) \geq d_{M \cap K_n[S]}(u_p) + 1 \geq \frac{k+2}{2}$, which contradicts to $\Delta(K_n[M]) \leq \frac{k}{2}$. Let us now assume $u_p \in S_2^1$. By the above procedure, there exists a vertex $u_q \in S_1^1$ such that when we pick up the edge $e_{1j} = u_pu_q$ $(1 \leq j \leq x_1)$ from $G_{1(j-1)}[S]$ the degree of u_p in $G_{1j}[S]$ is equal to $\frac{k-4}{2}$. That is $d_{G_{1j}[S]}(u_p) = \frac{k-4}{2}$ and $d_{G_{1(j-1)}[S]}(u_p) = \frac{k-2}{2}$. From our procedure, $|E_G[u_q, S_2^1]| = |E_{G_{1(j-1)}}[u_q, S_2^1]|$. Without loss of generality, let $|E_G[u_q, S_2^1]| = t$ and $u_qu_j \in E(G)$ for $x_1 + 1 \leq j \leq x_1 + t$; see Figure 3 (a). Thus $u_p \in \{u_{x_1+1}, u_{x_1+2}, \cdots, u_{x_1+t}\}$, and $u_qu_j \in M$ for $x_1 + t + 1 \leq j \leq k$. Because $|E_G[u_j, S_2^1]| \geq 1$ for each $u_j \in S_1^1$ $(1 \leq j \leq x_1)$, we have $t \geq 1$. Since $d_{G_{1(j-1)}[S]}(u_p) = \frac{k-2}{2}$, by our procedure $d_{G_{1(j-1)}[S]}(u_j) \leq \frac{k-2}{2}$ for each $u_j \in S_2^1$ $(x_1 + 1 \leq j \leq x_1 + t)$. Assume, to the contrary, that there is a vertex u_s $(x_1 + 1 \leq s \leq x_1 + t)$ such that $d_{G_{1(j-1)}[S]}(u_s) \geq \frac{k-2}{2}$. Then we should choose the edge u_qu_s instead of $e_{1j} = u_qu_p$ by our procedure, a contradiction. We conclude that $d_{G_{1(j-1)}[S]}(u_r) \leq \frac{k-2}{2}$ for each $u_r \in S_1^1$ $(x_1 + 1 \leq r \leq x_1 + t)$. Clearly, there are at least $k - 1 - \frac{k-2}{2} = \frac{k}{2}$ edges incident to each u_r $(x_1 + 1 \leq r \leq x_1 + t)$ belonging to $M \cup \{e_{11}, e_{12}, \cdots, e_{1(j-1)}\}$. Since $j \leq x_1$ and $u_qu_j \in M$ for $x_i + t + 1 \leq j \leq k$, we have $|E_{K_n[M]}[u_q, S_2^1]| + \sum_{j=1}^{t} d_{K_n[M]}(u_j) \geq k - x_1 - t + \frac{k}{2}t - (j-1) = k + \frac{k-2}{2}t - x_1 - j + 1$ and hence $|M| \geq |M \cap (E_{K_n}[w_1, S])| + \sum_{j=1}^{t} d_{K_n[M]}(u_j) + |E_{K_n[M]}[u_q, S_1^1]| \geq x_1 + (k + \frac{k-2}{2}t - x_1 - j + 1) = k + \frac{k-2}{2}t - j + 1$. Combining this with $j \leq x_1$, $t \geq 1$ and $x_1 \leq \frac{k}{2}$, we have $|M| \geq k + \frac{k}{2} - x_1 \geq k$, contradicting to $|M| = k - 1$.

By Claim 1, we have $\delta(G_1[S]) \geq \frac{k-2}{2}$. Recall that there exists at most one vertex in $K_n[M]$ such that its degree is $\frac{k}{2}$, and $x_{n-k} \leq x_{n-k-1} \leq \cdots \leq x_2 \leq x_1 \leq \frac{k}{2}$. Then $x_i \leq \frac{k-2}{2}$ for $2 \leq i \leq n - k$. Now we continue to introduce our procedure.

For $w_2 \in \bar{S}$, without loss of generality, let $S = S_1^2 \cup S_2^2$ and $S_1^2 = \{u_1, u_2, \cdots, u_{x_2}\}$ such that $u_jw_2 \in M$ for $1 \leq j \leq x_2$. Let $S_2^2 = S \setminus S_1^2 = \{u_{x_2+1}, u_{x_2+2}, \cdots, u_k\}$. Then $u_jw_2 \in E(G)$ for $x_2 + 1 \leq j \leq k$. Clearly, the tree $T_2' = w_2u_{x_2+1} \cup w_2u_{x_2+2} \cup \cdots \cup w_2u_k$ is a Steiner tree connecting S_2^2. Our idea is to seek for x_2 edges in $E_{G_1}[S_1^2, S_2^2]$ and add them to T_2' to form a Steiner tree connecting S. For each $u_j \in S_1^2$ $(1 \leq j \leq x_2)$, we



(a) (b) (c)

Figure 3. Graphs for Lemma 11.

claim that $|E_{G_1}[u_j, S_2^2]| \geq 1$. Otherwise, let $|E_{G_1}[u_j, S_2^2]| = 0$. Recall that $|M_1| = x_1$. Then there exist $k - x_2$ edges between u_j and S_2^2 belonging to $M \cup M_1$, and hence $|E_{K_n[M]}[u_j, S_2^2]| \geq k - x_2 - x_1$. Therefore, $|M| \geq |E_{K_n[M]}[u_j, S_2^2]| + d_{K_n[M]}(w_1) + d_{K_n[M]}(w_2) \geq (k - x_2 - x_1) + x_1 + x_2 = k$, which contradicts to $|M| = k - 1$. Choose the one with the smallest subscript among all the vertices of S_1^2 with the maximum degree in $G_1[S]$, say u_1'. Then search for the one adjacent to u_1' with the smallest subscript among all the vertices of S_2^2 with the maximum degree in $G_1[S]$, say u_1''. Let $e_{21} = u_1'u_1''$. Consider the graph $G_{21} = G_1 \setminus e_{21}$, and pick up the one with the smallest subscript among all the vertices of $S_1^2 \setminus u_1'$ with the maximum degree in $G_{21}[S]$, say u_2'. Then we search for the one adjacent to u_2' with the smallest subscript among all the vertices of S_2^2 with the maximum degree in $G_{21}[S]$, say u_2''. Set $e_{22} = u_2'u_2''$. Consider the graph $G_{22} = G_{21} \setminus e_{22} = G_1 \setminus \{e_{21}, e_{22}\}$. For each $u_j \in S_1^2$ $(1 \leq j \leq x_2)$, we proceed to find $e_{23}, e_{24}, \cdots, e_{2x_2}$ in the same way, and get graphs $G_{2j} = G_1 \setminus \{e_{21}, e_{22}, \cdots, e_{2(j-1)}\}$ $(1 \leq j \leq x_2)$. Let $M_2 = \{e_{21}, e_{22}, \cdots, e_{2x_2}\}$ and $G_2 = G_1 \setminus M_1$. Thus the tree $T_2 = w_2u_{x_2+1} \cup w_2u_{x_2+2} \cup \cdots \cup w_2u_k \cup e_{21} \cup e_{22} \cup \cdots \cup e_{2x_2}$ is our desired tree. Furthermore, T_2 and T_1 are two internally disjoint trees connecting S.

For $w_i \in \bar{S}$, without loss of generality, let $S = S_1^i \cup S_2^i$ and $S_1^i = \{u_1, u_2, \cdots, u_{x_i}\}$ such that $u_jw_i \in M$ for $1 \leq j \leq x_i$. Set $S_2^i = S \setminus S_1^i = \{u_{x_i+1}, u_{x_i+2}, \cdots, u_k\}$. Then $u_jw_i \in E(G)$ for $x_i + 1 \leq j \leq k$. One can see that the tree $T_i' = w_iu_{x_i+1} \cup w_iu_{x_i+2} \cup \cdots \cup w_iu_k$ is a Steiner tree connecting S_2^i. Our idea is to seek for x_i edges in $E_{G_{i-1}}[S_1^2, S_2^2]$ and add them to T_i' to form a Steiner tree connecting S. For each $u_j \in S_1^i$ $(1 \leq j \leq x_i)$, we claim that $|E_{G_{i-1}}[u_j, S_2^i]| \geq 1$. Otherwise, let $|E_{G_{i-1}}[u_j, S_2^i]| = 0$. Recall that $|M_j| = x_j$ $(1 \leq j \leq i)$. Then there are $k - x_i$ edges between u_j and S_2^i belonging to $M \cup (\bigcup_{j=1}^{i-1} M_j)$, and hence $|E_{K_n[M]}[u_j, S_2^i]| \geq k - x_i - \sum_{j=1}^{i-1} x_j$. Therefore, $|M| \geq |E_{K_n[M]}[u_j, S_2^i]| + \sum_{j=1}^{i} |M \cap (K_n[w_j, S])| \geq k - x_i - \sum_{j=1}^{i-1} x_j + \sum_{j=1}^{i} x_j = k$, contradicting to $|M| = k - 1$. Choose the one with the smallest subscript among all the vertices of S_1^i with the maximum degree in $G_{i-1}[S]$, say u_1'. Then search for the one adjacent to u_1' with the smallest subscript among all the vertices of S_2^i with the maximum degree in $G_{i-1}[S]$, say u_1''. Let $e_{i1} = u_1'u_1''$. Consider the graph $G_{i1} = G_{i-1} \setminus e_{i1}$, pick up the one with the smallest subscript among all the vertices of $S_1^i \setminus u_1'$ with the maximum degree in $G_{i1}[S]$, say u_2'. Then we search for the one adjacent to u_2' with the smallest subscript among all the vertices of S_2^i with the maximum degree in $G_{i1}[S]$, say u_2''. Set $e_{i2} = u_2'u_2''$. Consider the graph $G_{i2} = G_{i1} \setminus e_{i2} = G_{i-1} \setminus \{e_{i1}, e_{i2}\}$. For each $u_j \in S_1^i$ $(1 \leq j \leq x_i)$, we proceed to find $e_{i3}, e_{i4}, \cdots, e_{ix_i}$ in the same way, and get graphs $G_{ij} = G_{i-1} \setminus \{e_{i1}, e_{i2}, \cdots, e_{i(j-1)}\}$ $(1 \leq j \leq x_i)$. Let $M_i = \{e_{i1}, e_{i2}, \cdots, e_{ix_2}\}$ and $G_i = G_{i-1} \setminus M_i$. Thus the tree $T_i = w_iu_{x_2+1} \cup w_iu_{x_2+2} \cup \cdots \cup w_iu_k \cup e_{i1} \cup e_{i2} \cup \cdots \cup e_{ix_i}$ is our desired tree. Furthermore, $T_1, T_2, \cdots, T_i$ are pairwise internally disjoint trees connecting S.

We continue this procedure until we obtain $n-k$ pairwise internally disjoint trees $T_1, T_2, \cdots, T_{n-k}$. Note that if there exists some x_j such that $x_j = 0$ then $x_{j+1} = x_{j+2} = \cdots = x_{n-k} = 0$ since $x_1 \geq x_2 \geq \cdots \geq x_{n-k}$. Then $T_i = w_iu_1 \cup w_iu_2 \cup \cdots w_iu_k$ $(j \leq i \leq n-k)$ is our desired tree. From the above procedure, the resulting graph must be $G_{n-k} = G \setminus \bigcup_{i=1}^{n-k} M_i$. Let us show the following claim.

Claim 2. $\delta(G_{n-k}[S]) \geq \frac{k-2}{2}$.

Proof of Claim 2. Assume, to the contrary, that $\delta(G_{n-k}[S]) \leq \frac{k-4}{2}$, namely, there exists a vertex $u_p \in S$ such that $d_{G_{n-k}[S]}(u_p) \leq \frac{k-4}{2}$. Since $\delta(G[S]) \geq \frac{k-2}{2}$, by our procedure there exists an edge e_{ij} in $G_{i(j-1)}$ incident to the vertex u_p such that when we pick up this edge, $d_{G_{ij}[S]}(u_p) = \frac{k-4}{2}$ but $d_{G_{i(j-1)}[S]}(u_p) = \frac{k-2}{2}$.

First, we consider the case $u_p \in S_2^i$. Then there exists a vertex $u_q \in S_1^i$ such that when we pick up the edge $e_{ij} = u_pu_q$ from $G_{i(j-1)}[S]$ the degree of u_p in $G_{ij}[S]$ is equal to $\frac{k-4}{2}$, that is, $d_{G_{ij}[S]}(u_p) = \frac{k-4}{2}$ and $d_{G_{i(j-1)}[S]}(u_p) = \frac{k-2}{2}$. From our procedure, $|E_{G_{i-1}}[u_q, S_2^i]| = |E_{G_{i(j-1)}}[u_q, S_2^i]|$. Without loss of generality, let $|E_{G_{i-1}}[u_q, S_2^i]| = t$ and $u_qu_j \in E(G_{i-1})$ for $x_i+1 \leq j \leq x_i+t$; see Figure 3 (b). Thus $u_p \in \{u_{x_i+1}, u_{x_i+2}, \cdots, u_{x_i+t}\}$, and $u_qu_j \in M \cup (\bigcup_{r=1}^{i-1} M_r)$ for $x_i+t+1 \leq j \leq k$. Since $x_i \leq \frac{k-2}{2}$ $(2 \leq i \leq n-k)$, it follows that $|S_1^i| \leq \frac{k-2}{2}$. From this together with $\delta(G_{i-1}[S]) \geq \frac{k-2}{2}$, we have $|E_{G_{i-1}}[u_q, S_1^i]| \geq 1$, that is, $t \geq 1$. Since $d_{G_{i(j-1)}[S]}(u_p) = \frac{k-2}{2}$, by our procedure $d_{G_{i(j-1)}[S]}(u_j) \leq \frac{k-2}{2}$ for each $u_j \in S_2^i$ $(x_i+1 \leq j \leq x_i+t)$. Assume, to the contrary, that there exists a vertex u_s $(x_i+1 \leq s \leq x_i+t)$ such that $d_{G_{i(j-1)}[S]}(u_s) \geq \frac{k-2}{2}$. Then we should choose the edge u_qu_s instead of $e_{ij} = u_qu_p$ by our procedure, a contradiction. We conclude that $d_{G_{i(j-1)}[S]}(u_r) \leq \frac{k-2}{2}$ for each $u_r \in S_2^i$ $(x_i+1 \leq r \leq x_i+t)$. Clearly, there are at least $k-1-\frac{k-2}{2} = \frac{k}{2}$ edges incident to each u_r $(x_i+1 \leq r \leq x_i+t)$ belonging to $M \cup (\bigcup_{j=1}^{i-1} M_j) \bigcup \{e_{i1}, e_{i2}, \cdots, e_{i(j-1)}\}$. Since $j \leq x_i$ and $u_qu_j \in M \cup (\bigcup_{r=1}^{i-1} M_r)$ for $x_i+t+1 \leq j \leq k$, we have

$$\begin{aligned}
|E_{K_n[M]}[u_q, S_2^i]| + \sum_{j=1}^{t} d_{K_n[M]}(u_j) &\geq k - x_i - t + \frac{k}{2}t - \sum_{j=1}^{i-1} x_j - (j-1) \\
&\geq k + \frac{k-2}{2}t - \sum_{j=1}^{i} x_j - x_i + 1 \quad (\textit{since } j \leq x_i)
\end{aligned}$$

and hence

$$\begin{aligned} |M| &\geq \sum_{j=1}^{i} |M \cap (E_{K_n}[w_j, S])| + \sum_{j=1}^{t} d_{K_n[M]}(u_j) + |E_{K_n[M]}[u_q, S_2^i]| \\ &\geq \sum_{j=1}^{i} x_j + k + \frac{k-2}{2}t - \sum_{j=1}^{i} x_j - x_i + 1 \\ &= k + \frac{k-2}{2}t - x_i + 1 \\ &\geq k + \frac{k-2}{2} - x_i + 1 && (\textit{since } t \geq 1) \\ &\geq k+1, && \left(\textit{since } x_i \leq \frac{k-2}{2}\right) \end{aligned}$$

which contradicts to $|M| = k-1$.

Next, assume $u_p \in S_1^i$. Recall that $d_{G_{ij}[S]}(u_p) = \frac{k-4}{2}$. Since $u_p \in S_1^i$, it follows that $d_{G_{i-1}[S]}(u_p) = \frac{k-2}{2}$. If $u_p \in \bigcap_{j=1}^{i} S_1^j$, namely, $u_p w_j \in M$ $(1 \leq j \leq i)$, then by our procedure $d_{G[S]}(u_p) = \frac{k-2}{2} + i - 1$ and hence $d_{K_n[S] \cap M}(u_p) = k - 1 - (\frac{k-2}{2} + i - 1) = \frac{k}{2} - i + 1$. Since $u_p w_j \in M$ for each $w_j \in \bar{S}$ $(1 \leq j \leq i)$, we have $d_{K_n[M]}(u_p) = d_{K_n[S] \cap M}(u_p) + d_{K_n[S,\bar{S}] \cap M}(u_p) \geq (\frac{k}{2} - i + 1) + i = \frac{k+2}{2}$, contradicting to $\Delta(K_n[M]) \leq \frac{k}{2}$. Combining this with $u_p \in S_1^i$, we have $u_p \notin \bigcap_{j=1}^{i-1} S_1^i$ and we can assume that there exists an integer i' $(i' \leq i-1)$ satisfying the following conditions:

- $u_p \in S_2^{i'}$ and $d_{G_{i'}[S]}(u_p) < d_{G_{i'-1}[S]}(u_p)$;
- if u_p belongs to some S_2^j $(i'+1 \leq j \leq i)$ then $d_{G_j[S]}(u_p) = d_{G_{j-1}[S]}(u_p)$.

The above two conditions can be restated as follows:

- $u_p w_{i'} \in E(G)$ and $d_{G_{i'}[S]}(u_p) < d_{G_{i'-1}[S]}(u_p)$;
- if $u_p w_j \in E(G)$ $(i'+1 \leq j \leq i)$ then $d_{G_j[S]}(u_p) = d_{G_{j-1}[S]}(u_p)$.

In fact, we can find the integer i' such that $u_p w_{i'} \in E(G)$ and $d_{G_{i'}[S]}(u_p) < d_{G_{i'-1}[S]}(u_p)$. Assume, to the contrary, that for each w_j $(1 \leq j \leq i)$, $u_p w_j \in M$, or $u_p w_j \in E(G)$ but $d_{G_j[S]}(u_p) = d_{G_{j-1}[S]}(u_p)$. Let i_1 $(i_1 \leq i)$ be the number of vertices nonadjacent to $u_p \in S$ in $\{w_1, w_2, \cdots, w_{i-1}\} \subseteq \bar{S}$. Without loss of generality, let $w_j u_p \in M$ $(1 \leq j \leq i_1)$. Recall that $d_{G_{ij}[S]}(u_p) = \frac{k-4}{2}$. Thus $d_{G[S]}(u_p) = \frac{k-4}{2} + i_1$ and hence $d_{K_n[S] \cap M}(u_p) \geq k - 1 - (\frac{k-4}{2} + i_1) = \frac{k+2}{2} - i_1$. Since $w_j u_p \in M$ $(1 \leq j \leq i_1)$, it follows that $d_{K_n[S,\bar{S}] \cap M}(u_p) \geq i_1$, which results in $d_{K_n[M]}(u_p) = d_{K_n[S] \cap M}(u_p) + d_{K_n[S,\bar{S}] \cap M}(u_p) \geq (\frac{k+2}{2} - i_1) + i_1 = \frac{k+2}{2}$, contradicting to $\Delta(K_n[M]) \leq \frac{k}{2}$.

Now we turn our attention to $u_p \in S_2^{i'}$. Without loss of generality, let $u_p w_j \in M$ $(j \in \{j_1, j_2, \cdots, j_{i_1}\})$, namely, $u_p \in S_1^{j_1} \cap S_1^{j_2} \cap \cdots \cap S_1^{j_{i_1}}$, where $j_1, j_2, \cdots, j_{i_1} \in \{i'+1, i'+2, \cdots, i\}$. Then $u_p w_j \in E(G)$ $(j \in \{i'+1, i'+2, \cdots, i\} \setminus \{j_1, j_2, \cdots, j_{i_1}\})$. Clearly, $i_1 \leq i - i'$. Recall that $u_p \in S_1^i$ and $d_{G_{ij}[S]}(u_p) = \frac{k-4}{2}$. Thus $d_{G_{i'}[S]}(u_p) = \frac{k-4}{2} + i_1$. By our procedure, there exists a vertex $u_q \in S_1^{i'}$ such that when we pick up the edge $e_{i'j} = u_p u_q$ from $G_{i'(j-1)}[S]$ the degree of u_p in $G_{i'j}[S]$ is equal to $\frac{k-4}{2} + i_1$, that is, $d_{G_{i'j}[S]}(u_p) = \frac{k-4}{2} + i_1$ and $d_{G_{i'(j-1)}[S]}(u_p) = \frac{k-2}{2} + i_1$. From our procedure, $|E_{G_{i'-1}}[u_q, S_2^{i'}]| = |E_{G_{i'(j-1)}}[u_q, S_2^{i'}]|$. Without loss of generality, let $|E_{G_{i'-1}}[u_q, S_2^{i'}]| = t$ and $u_qu_j \in E(G_{i'-1})$ for $x_{i'}+1 \le j \le x_{i'}+t$; see Figure 3 (c). Thus $u_p \in \{u_{x_{i'}+1}, u_{x_{i'}+2}, \cdots, u_{x_{i'}+t}\}$, and $u_qu_j \in M \cup (\bigcup_{r=1}^{i'-1} M_r)$ for $x_{i'}+t+1 \le j \le k$. Since $x_j \le \frac{k-2}{2}$ $(2 \le j \le n-k)$, it follows that $|S_1^{i'}| \le \frac{k-2}{2}$. From this together with $\delta(G_{i'-1}[S]) \ge \frac{k-2}{2}$, we have $|E_{G_{i'-1}}[u_q, S_1^{i'}]| \ge 1$, that is, $t \ge 1$. Since $d_{G_{i'(j-1)}[S]}(u_p) = \frac{k-2}{2}+i_1$, by our procedure $d_{G_{i'(j-1)}[S]}(u_j) \le \frac{k-2}{2}+i_1$ for each $u_j \in S_2^{i'}$ $(x_{i'}+1 \le j \le x_{i'}+t)$. Assume, to the contrary, that there is a vertex u_s $(x_{i'}+1 \le s \le x_{i'}+t)$ such that $d_{G_{i'(j-1)}[S]}(u_s) \ge \frac{k-2}{2}+i_1+1$. Then we should choose the edge u_qu_s instead of $e_{i'j} = u_qu_p$ by our procedure, a contradiction. We conclude that $d_{G_{i'(j-1)}[S]}(u_r) \le \frac{k-2}{2}+i_1$ for each $u_r \in S_2^{i'}$ $(x_{i'}+1 \le r \le x_{i'}+t)$. Clearly, there are at least $k-1-(\frac{k-2}{2}+i_1) = \frac{k}{2}-i_1$ edges incident to each u_r $(x_{i'}+1 \le r \le x_{i'}+t)$ belonging to $M \cup (\bigcup_{j=1}^{i'-1} M_j) \bigcup \{e_{i'1}, e_{i'2}, \cdots, e_{i'(j-1)}\}$. Since $j \le x_{i'}$ and $u_qu_j \in M \cup (\bigcup_{r=1}^{i'-1} M_r)$ for $x_{i'}+t+1 \le j \le k$, we have

$$
\begin{aligned}
& |E_{K_n[M]}[u_q, S_2^{i'}]| + \sum_{j=1}^{t} d_{K_n[M]}(u_j) \\
\geq\ & k - x_{i'} - t + \left(\frac{k}{2} - i_1\right) t - \sum_{j=1}^{i'-1} x_j - (j-1) \\
\geq\ & k - \sum_{j=1}^{i'} x_j + \left(\frac{k-2}{2} - i_1\right) t - x_{i'} + 1 \qquad (\textit{since } j \le x_{i'}) \\
\geq\ & k - \sum_{j=1}^{i'} x_j + \left(\frac{k-2}{2} - i + i'\right) t - x_{i'} + 1 \quad (\textit{since } i_1 \le i - i')
\end{aligned}
$$

and hence

$$
\begin{aligned}
|M| &\geq \sum_{j=1}^{i} |M \cap (E_{K_n}[w_j, S])| + \sum_{j=1}^{p} d_{K_n[M]}(u_j) + |E_{K_n[M]}[u_q, S_2^{i}]| \\
&\geq \sum_{j=1}^{i} x_j + k - \sum_{j=1}^{i'} x_j + \left(\frac{k-2}{2} - i + i'\right) t - x_{i'} + 1 \\
&= k + \sum_{j=i'+1}^{i} x_j + \frac{k-2}{2} - (i - i') - x_{i'} + 1 \ (\textit{since } t \ge 1) \\
&\geq k + 1, \qquad \left(\textit{since } x_{i'} \le \frac{k-2}{2} \textit{ and } x_j \ge 1 \textit{ for } i'+1 \le j \le i\right)
\end{aligned}
$$

contradicting $|M| = k-1$. This completes the proof of Claim 2.

From our procedure, we get $n-k$ internally disjoint trees connecting S in G, say $T_1, T_2, \cdots, T_{n-k}$. Recall that $G_{n-k} = G \setminus \bigcup_{i=1}^{n-k} M_i$. We can regard $G_{n-k}[S] = G[S] \setminus (\bigcup_{i=1}^{n-k} M_i)$ as a graph obtained from the complete graph K_k by deleting $|M'| + \sum_{i=1}^{n-k} |M_i|$ edges. Since $|M'| + \sum_{i=1}^{n-k} |M_i| + |M''| = m_1 + \sum_{i=1}^{n-k} x_i + m_2 = k-1$, we have $1 \le \sum_{i=1}^{n-k} |M_i| + m_1 \le k-1$. By Claim 2, $\delta(G_{n-k}[S]) \ge \frac{k-2}{2}$ and hence $2 \le \Delta(\overline{G_{n-k}}[S]) \le \frac{k}{2}$. From Lemma 9, there exist $\frac{k-2}{2}$ edge-disjoint spanning trees connecting S in $G_{n-k}[S]$. These trees together with $T_1, T_2, \cdots, T_{n-k}$ are $n - \frac{k}{2} - 1$ internally disjoint trees connecting S in G, namely, $\kappa(S) \ge n - \frac{k}{2} - 1$. From the arbitrariness of S, we have $\kappa_k(G) \ge n - \frac{k}{2} - 1$, as desired. $\square$

We can now prove our main results.

Proof of Theorem 2. Assume that $\kappa_k(G) = n - \frac{k}{2} - 1$. Since G of order n is connected, we can regard G as a graph obtained from the complete graph K_n by deleting some edges. From Lemma 4, it follows that $|M| \ge 1$ and hence $\Delta(K_n[M]) \ge 1$. If $G = K_n \setminus M$ where $M \subseteq E(K_n)$ such that $\Delta(K_n[M]) \ge \frac{k}{2} + 1$, then $\kappa_k(G) \le \lambda_k(G) < n - \frac{k}{2} - 1$ by Observation 1 and Corollary 1, a contradiction. So $1 \le \Delta(K_n[M]) \le \frac{k}{2}$. If $2 \le \Delta(K_n[M]) \le \frac{k}{2}$ and $|M| \ge k$, then $\kappa_k(G) \le \lambda_k(G) < n - \frac{k}{2} - 1$ by Observation 1 and Lemma 7, a contradiction. Therefore, $1 \le |M| \le k-1$. If $\Delta(K_n[M]) = 1$, then $1 \le |M| \le k-1$ by Lemma 8. We conclude that $1 \le \Delta(K_n[M]) \le \frac{k}{2}$ and $1 \le |M| \le k-1$, as desired.

Conversely, let $G = K_n \setminus M$ where $M \subseteq E(K_n)$ such that $1 \le \Delta(K_n[M]) \le \frac{k}{2}$ and $1 \le |M| \le k-1$. In fact, we only need to show that $\kappa_k(G) \ge n - \frac{k}{2} - 1$ for $\Delta(K_n[M]) = 1$ and $|M| = k-1$, or $2 \le \Delta(K_n[M]) \le \frac{k}{2}$ and $|M| = k-1$. The results follow by (1) of Lemma 10 and Lemma 11.

Proof of Theorem 3. If G is a connected graph satisfying the condition (2), then $\kappa_k(G) = n - \frac{k}{2} - 1$ by Theorem 2. From Observation 1, $\lambda_k(G) \ge \kappa_k(G) = n - \frac{k}{2} - 1$. Combining this with Lemma 4, we have $\lambda_k(G) = n - \frac{k}{2} - 1$. Assume that G is a connected graph satisfying the condition (1). We only need to show that $\lambda_k(G) = n - \frac{k}{2} - 1$ for $|M| = \lfloor \frac{n}{2} \rfloor$. The result follows by (2) of Lemma 10 and Lemma 5.

Conversely, assume that $\lambda_k(G) = n - \frac{k}{2} - 1$. Since G of order n is connected, we can consider G as a graph obtained from a complete graph K_n by deleting some edges. From Corollary 1, $G = K_n \setminus M$ such that $\Delta(K_n[M]) \le \frac{k}{2}$, where $M \subseteq E(K_n)$. Combining this with Lemma 5, we have $|M| \ge 1$ and $\Delta(K_n[M]) \ge 1$. So $1 \le \Delta(K_n[M]) \le \frac{k}{2}$. It is clear that if $\Delta(K_n[M]) = 1$ then $1 \le |M| \le \lfloor \frac{n}{2} \rfloor$. If $2 \le \Delta(K_n[M]) \le \frac{k}{2}$, then $1 \le |M| \le k-1$ by Lemma 7. So (1) or (2) holds.

Remark 3. As we know, $\lambda(G) = n-2$ if and only if $G = K_n \setminus M$ such that $\Delta(K_n[M]) = 1$ and $1 \le |M| \le \lfloor \frac{n}{2} \rfloor$, where $M \subseteq E(K_n)$. So we can restated the above conclusion as follows: $\lambda_2(G) = n-2$ if and only if $G = K_n \setminus M$ such that $\Delta(K_n[M]) = 1$ and $1 \le |M| \le \lfloor \frac{n}{2} \rfloor$, where $M \subseteq E(K_n)$. This means that $4 \le k \le n$ in Theorem 3 can be replaced by $2 \le k \le n$.

References

[1] J.A. Bondy, U.S.R. Murty, *Graph Theory*, GTM 244, Springer, 2008.

[2] G. Chartrand, S.F. Kappor, L. Lesniak, D.R. Lick, *Generalized connectivity in graphs*, Bull. Bombay Math. Colloq. 2(1984), 1-6.

[3] G. Chartrand, F. Okamoto, P. Zhang, *Rainbow trees in graphs and generalized connectivity*, Networks 55(4)(2010), 360-367.

[4] M. Grötschel, *The Steiner tree packing problem in $VLSI$ design*, Math. Program. 78(1997), 265-281.

[5] M. Grötschel, A. Martin, R. Weismantel, *Packing Steiner trees: A cutting plane algorithm and commputational results*, Math. Program. 72(1996), 125-145.

[6] K. Jain, M. Mahdian, M. Salavatipour, *Packing Steiner trees*, in: Pro. of the 14th ACM-$SIAM$ symposium on Discterte Algorithms, Baltimore, 2003, 266-274.

[7] M. Kriesell, *Edge-disjoint trees containing some given vertices in a graph*, J. Combin. Theory, Ser.B 88(2003), 53-65.

[8] M. Kriesell, *Edge-disjoint Steiner trees in graphs without large bridges*, J. Graph Theory 62(2009), 188-198.

[9] L. Lau, *An approxinate max-Steiner-tree-packing min-Steiner-cut theorem*, Combinatorica 27(2007), 71-90.

[10] H. Li, X. Li, Y. Sun, *The generalied* 3*-connectivity of Cartesian product graphs*, Discrete Math. Theor. Comput. Sci. 14(1)(2012), 43-54.

[11] S. Li, W. Li, X. Li, *The generalized connectivity of complete bipartite graphs*, Ars Combin. 104(2012), 65-79.

[12] S. Li, W. Li, X. Li, *The generalized connectivity of complete equipartition* 3*-partite graphs*, Bull. Malays. Math. Sci. Soc., in press.

[13] S. Li, X. Li, *Note on the hardness of generalized connectivity*, J. Combin. Optimization 24(2012), 389-396.

[14] S. Li, X. Li, W. Zhou, *Sharp bounds for the generalized connectivity $\kappa_3(G)$*, Discrete Math. 310(2010), 2147-2163.

[15] X. Li, Y. Mao, Y. Sun, *On the generalized (edge-)connectivity of graphs*, arXiv:1112.0127 [math.CO] 2011.

[16] C.St.J.A. Nash-Williams, *Edge-disjonint spanning trees of finite graphs*, J. London Math. Soc. 36(1961), 445-450.

[17] F. Okamoto, P. Zhang, *The tree connectivity of regular complete bipartite graphs*, J. Combin. Math. Combin. Comput. 74(2010), 279-293.

[18] N.A. Sherwani, *Algorithms for VLSI Physical Design Automation*, 3rd Edition, Kluwer Acad. Pub., London, 1999.

[19] W. Tutte, *On the problem of decomposing a graph into n connected factors*, J. London Math. Soc. 36(1961), 221-230.

[20] D. West, H. Wu, *Packing Steiner trees and S-connectors in graphs*, J. Combin. Theory, Ser.B 102(2012), 186-205.

Appendix

An example for Case 2 of Lemma 11

Let $k = 8$, and $G = K_n \setminus M$ where $M \subseteq E(K_n)$ be a connected graph of order n such that $|M| = k - 1 = 7$ and $\Delta(K_n[M]) \leq \frac{k}{2} = 4$. Let $S = \{u_1, u_2, \cdots, u_8\}$, $\bar{S} = V(G) \setminus S = \{w_1, w_2, \cdots, w_{n-8}\}$ and $M = \{w_1u_1, w_1u_2, w_1u_3, w_2u_2, w_2u_4, u_5u_6, u_6u_8\}$; see Figure 4 (a). Clearly, $x_1 = |E_{K_n[M]}[w_1, S]| = 3 \geq x_2 = |E_{K_n[M]}[w_2, S]| = 2 > x_i = |E_{K_n[M]}[w_i, S]| = 0$ $(3 \leq i \leq n - 8)$.

For w_1, we let $S_1^1 = \{u_1, u_2, u_3\}$ since $w_1u_1, w_1u_2, w_1u_3 \in M$. Set $S_2^1 = S \setminus S_1^1 = \{u_4, u_5, u_6, u_7, u_8\}$. Clearly, $d_{G[S]}(u_1) = d_{G[S]}(u_2) = d_{G[S]}(u_3) = 7 = k - 1$ and hence u_1, u_2, u_3 are all the vertices of S_1^1 with the maximum degree in $G[S]$. But u_1 is the one with the smallest subscript, so we choose $u_1' = u_1$ in S_1^1 and search for the vertex adjacent to u_1' in S_2^1 and obtain $u_4, u_5, u_6, u_7, u_8 \in S_2^1$ since $u_1'u_j \in E(G)$ $(j = 4, \cdots, 8)$. Obviously, $d_{G[S]}(u_4) = d_{G[S]}(u_7) = 7 > d_{G[S]}(u_5) = d_{G[S]}(u_8) = 6 > d_{G[S]}(u_6) = 5$ and hence u_4, u_7 are two vertices of S_2^1 with the maximum degree in $G[S]$. Since u_4 is the one with the smallest subscript, we pick up $u_1'' = u_4 \in S_2^1$ and put $e_{11} = u_1'u_1''(= u_1u_4)$. Consider the graph $G_{11} = G \setminus e_{11}$. Since $d_{G_{11}[S]}(u_2) = d_{G_{11}[S]}(u_3) = 7$ and the subscript of u_2 is smaller than u_3, we choose $u_2' = u_2$ in $S_1^1 \setminus u_1'$ and search for the vertices adjacent to u_2' in S_2^1 and obtain $u_4, u_5, u_6, u_7, u_8 \in S_2^1$ since $u_2'u_j \in E(G_{11})$ $(j = 4, \cdots, 8)$. Since $d_{G_{11}[S]}(u_7) = 7 > d_{G_{11}[S]}(u_j) = 6 > d_{G_{11}[S]}(u_6) = 5$ $(j = 4, 5, 8)$, we pick up $u_2'' = u_7 \in S_2^1$ and get $e_{12} = u_2'u_2''$ $(= u_2u_7)$. Consider the graph $G_{12} = G_{11} \setminus e_{12} = G \setminus \{e_{11}, e_{12}\}$. There is only one vertex u_3 in $S_1 \setminus \{u_1', u_2'\} = S_1 \setminus \{u_1, u_2\}$. Therefore, let $u_3' = u_3$ and search for the vertices adjacent to u_3' in S_2^1 and obtain $u_j \in S_2^1$ since $u_3'u_j \in E(G_{12})$ $(j = 4, \cdots, 8)$. Since $d_{G_{12}[S]}(u_j) = 6 > d_{G_{12}[S]}(u_6) = 5$ $(i = 4, 5, 7, 8)$, it follows that u_4, u_5, u_7, u_8 are all the vertices of S_2^1 with the maximum degree in $G_{12}[S]$. But u_4 is the one with the smallest subscript, so we choose $u_3'' = u_4 \in S_2^1$ and get $e_{13} = u_3'u_3''$ $(= u_3u_4)$. Since $x_1 = |E_{K_n[M]}[w_1, S]| = 3$, we terminate this procedure. Set $M_1 = \{e_{11}, e_{12}, e_{13}\}$ and $G_1 = G \setminus M_1$. Thus the tree $T_1 = w_1u_4 \cup w_1u_5 \cup w_1u_6 \cup w_1u_7 \cup w_1u_8 \cup u_1u_4 \cup u_2u_7 \cup u_3u_4$ is our desired tree; see Figure 4 (b).

For w_2, we let $S_1^2 = \{u_2, u_4\}$ since $w_2u_2, w_2u_4 \in M$. Let $S_2^2 = S \setminus S_1^2 = \{u_1, u_3, u_5, u_6, u_7, u_8\}$. Since $d_{G_1[S]}(u_2) = 6 > d_{G_1[S]}(u_4) = 5$, it follows that u_2 is the vertex of S_1^2 with the maximum degree in $G_1[S]$. So we choose $u_1' = u_2$ in S_1^2 and search for the vertices adjacent to u_1' $(= u_2)$ in S_2^2 and obtain $u_1, u_3, u_5, u_6, u_8 \in S_2^2$ since $u_1'u_j \in E(G_{21})$ $(j = 1, 3, 5, 6, 8)$. Since $d_{G_1[S]}(u_j) = 6 > d_{G_1[S]}(u_6) = 5$ $(j = 1, 3, 5, 8)$ and u_1 is the vertex of the maximum degree with the smallest subscript, we choose $u_1'' = u_1 \in S_2^2$. Put $e_{21} = u_1'u_1''$ $(= u_2u_1)$. Consider the graph $G_{21} = G_1 \setminus e_{21}$. Clearly, $S_1 \setminus \{u_1'\} = S_1 \setminus \{u_2\} = \{u_4\}$, so we let $u_2' = u_4$ and search for the vertices adjacent to u_2' $(= u_4)$ in S_2^2 and obtain u_5, u_6, u_7, u_8 since $u_2u_j \in E(G)$ $(j = 5, 6, 7, 8)$. Since $d_{G_{21}[S]}(u_j) = 6 > d_{G_{21}[S]}(u_6) = 5$ $(j = 5, 7, 8)$ and u_5 is the one with the smallest subscript, we choose $u_2'' = u_5 \in S_2^2$ and get $e_{22} = u_2'u_2''$ $(= u_4u_5)$. Since $x_2 = |E_{K_n[M]}[w_2, S]| = 2$, we terminate this procedure. Let $M_2 = \{e_{21}, e_{22}\}$ and



(a) (b) (c)

(d) (e)

Figure 4 Graphs for the appendix.

$G_2 = G_1 \backslash M_2$. Then the tree $T_2 = w_2u_1 \cup w_2u_3 \cup w_2u_5 \cup w_2u_6 \cup w_2u_7 \cup w_2u_8 \cup u_2u_1 \cup u_4u_5$ is our desired tree; see Figure 4 (c). Obviously, T_2 and T_1 are two internally disjoint trees connecting S.

Since $x_i = |E_{K_n[M]}[w_i, S]| = 0$ for $3 \leq i \leq n-8$, we terminate this procedure. For $w_3, \cdots, w_{n-8}$, the trees $T_i = w_iu_1 \cup w_iu_2 \cup \cdots \cup w_iu_8$ $(3 \leq i \leq n-8)$ (see Figure 4 (d)) are our desired trees.

We can consider $G_2[S] = G[S] \setminus \{M_1, M_2\}$ as a graph obtained from complete graph K_k by deleting $|M \cap K_n[S]| + |M_1| + |M_2|$ edges. Since $|M \cap K_n[S]| + |M_1| + |M_2| = 2 + 3 + 2 = 7 = k - 1$, from Lemma 9, there exist 3 edge-disjoint spanning trees connecting S in $G[S]$ (Actually, we can give 3 edge-disjoint trees spanning connecting S; see Figure 4 (e). For example, the trees $T_1' = u_1u_8 \cup u_8u_4 \cup u_4u_6 \cup u_6u_3 \cup u_3u_2 \cup u_2u_5 \cup u_5u_7$, $T_2' = u_4u_7 \cup u_7u_8 \cup u_8u_3 \cup u_3u_1 \cup u_1u_5 \cup u_1u_6 \cup u_6u_2$ and $T_3' = u_2u_4 \cup u_2u_8 \cup u_8u_5 \cup u_5u_3 \cup u_3u_7 \cup u_1u_7 \cup u_7u_6$ can be our desired trees). These three trees together with $T_1, T_2, \cdots, T_{n-8}$ are $n - 5 = n - \frac{k}{2} - 1$ internally disjoint trees connecting S, namely, $\lambda(S) \geq n - 5$.